










Radiation Dose to Patients From Common Imaging Examinations

Procedure		**Approximate effective radiation dose	Comparable to natural background radiation for	* Estimated lifetime risk of fatal cancer from examination
 ABDOMINAL REGION	Computed Tomography (CT) — Abdomen and Pelvis	10 mSv	3 years	Low
	Computed Tomography (CT) — Abdomen and Pelvis, repeated with and without contrast material	20 mSv	7 years	Moderate
	Computed Tomography (CT) — Colonography	10 mSv	3 years	Low
	Intravenous Pyelogram (IVP)	3 mSv	1 year	Low
	Radiography (X-ray) — Lower GI Tract	8 mSv	3 years	Low
	Radiography (X-ray) — Upper GI Tract	6 mSv	2 years	Low
 BONE	Radiography (X-ray) — Spine	1.5 mSv	6 months	Very Low
	Radiography (X-ray) — Extremity	0.001 mSv	3 hours	Negligible
 CENTRAL NERVOUS SYSTEM	Computed Tomography (CT) — Head	2 mSv	8 months	Very Low
	Computed Tomography (CT) — Head, repeated with and without contrast material	4 mSv	16 months	Low
	Computed Tomography (CT) — Spine	6 mSv	2 years	Low
 CHEST	Computed Tomography (CT) — Chest	7 mSv	2 years	Low
	Computed Tomography (CT) — Lung Cancer Screening	1.5 mSv	6 months	Very Low
	Radiography — Chest	0.1 mSv	10 days	Minimal
 DENTAL	Intraoral X-ray	0.005 mSv	1 day	Negligible
 HEART	Coronary Computed Tomography Angiography (CTA)	12 mSv	4 years	Low
	Cardiac CT for Calcium Scoring	3 mSv	1 year	Low
 MEN'S IMAGING	Bone Densitometry (DEXA)	0.001 mSv	3 hours	Negligible
 NUCLEAR MEDICINE	Positron Emission Tomography — Computed Tomography (PET/CT)	25 mSv	8 years	Moderate
 WOMEN'S IMAGING	Bone Densitometry (DEXA)	0.001 mSv	3 hours	Negligible
	Mammography	0.4 mSv	7 weeks	Very Low

*Risk Level	Negligible	Minimal	Very Low	Low	Moderate
Estimated additional risk of fatal cancer for an adult from examination	Less than 1 in 1,000,000	1 in 1,000,000 to 1 in 100,000	1 in 100,000 to 1 in 10,000	1 in 10,000 to 1 in 1,000	1 in 1,000 to 1 in 500
Note: These risk levels represent very small additions to the 1 in 5 chance we all have of dying from cancer.					

Important: Pediatric patients vary in size. Doses given to pediatric patients will vary significantly from those given to adults.

RadiologyInfo.org
The radiology information resource for patients

RSNA
Radiological Society
of North America

ACR
AMERICAN COLLEGE OF
RADIOLOGY
QUALITY IS OUR IMAGE

For the most current information, visit radiologyinfo.org.

** The effective doses are typical values for an average-sized adult. The actual dose can vary substantially, depending on a person's size as well as on differences in imaging practices.†

†Mettler, F.A., et al. "Effective doses in radiology and diagnostic nuclear medicine: a catalog." *Radiology*, July 2008; 248(1):254–263.